

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC207708L3V

MBD1 (NM_015844) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	MBD1 (NM_015844) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MBD1
Synonyms:	CXXC3; PCM1; RFT
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_015844
ORF Size:	1647 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC207708).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 015844.1</u>
RefSeq Size:	2793 bp
RefSeq ORF:	1650 bp
Locus ID:	4152
UniProt ID:	<u>Q9UIS9</u>
Cytogenetics:	18q21.1
Protein Families:	Druggable Genome, Transcription Factors
MW:	59.8 kDa



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:The protein encoded by this gene is a member of a family of nuclear proteins related by the
presence of a methyl-CpG binding domain (MBD). These proteins are capable of binding
specifically to methylated DNA, and some members can also repress transcription from
methylated gene promoters. This protein contains multiple domains: MBD at the N-terminus
that functions both in binding to methylated DNA and in protein interactions; several CXXC-
type zinc finger domains that mediate binding to non-methylated CpG dinucleotides;
transcriptional repression domain (TRD) at the C-terminus that is involved in transcription
repression and in protein interactions. Numerous alternatively spliced transcript variants
encoding different isoforms have been noted for this gene.[provided by RefSeq, Feb 2011]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US