

## Product datasheet for RC207721L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TAB2 (NM\_015093) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** TAB2 (NM\_015093) Human Tagged ORF Clone Lentiviral Particle

Symbol: TAB2

**Synonyms:** CHTD2; MAP3K7IP2; TAB-2

**Mammalian Cell** 

Selection:

ACCN:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 015093

Tag: Myc-DDK

ORF Size: 2079 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC207721).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 015093.2</u>

 RefSeq Size:
 4414 bp

 RefSeq ORF:
 2082 bp

 Locus ID:
 23118

 UniProt ID:
 Q9NYJ8

 Cytogenetics:
 6q25.1

**Domains:** zf-RanBP, CUE

**Protein Families:** Druggable Genome





## TAB2 (NM\_015093) Human Tagged ORF Clone Lentiviral Particle - RC207721L3V

Protein Pathways: MAPK signaling pathway, NOD-like receptor signaling pathway, Toll-like receptor signaling

pathway

**MW:** 76.5 kDa

**Gene Summary:** The protein encoded by this gene is an activator of MAP3K7/TAK1, which is required for for

the IL-1 induced activation of nuclear factor kappaB and MAPK8/JNK. This protein forms a kinase complex with TRAF6, MAP3K7 and TAB1, and it thus serves as an adaptor that links MAP3K7 and TRAF6. This protein, along with TAB1 and MAP3K7, also participates in the signal transduction induced by TNFSF11/RANKI through the activation of the receptor activator of NF-kappaB (TNFRSF11A/RANK), which may regulate the development and function of osteoclasts. Studies of the related mouse protein indicate that it functions to protect against liver damage caused by chemical stressors. Mutations in this gene cause congenital heart defects, multiple types, 2 (CHTD2). Alternative splicing results in multiple transcript variants.

[provided by RefSeq, May 2014]