

## Product datasheet for RC207731L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **CLCA2 (NM\_006536) Human Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

**Product Type:** Lentiviral Particles

Product Name: CLCA2 (NM 006536) Human Tagged ORF Clone Lentiviral Particle

Symbol: CLCA2

**Synonyms:** CACC; CaCC-3; CACC3; CLCRG2

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag:Myc-DDKACCN:NM\_006536

ORF Size: 2829 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC207731).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 006536.4

RefSeq Size: 4043 bp
RefSeq ORF: 2832 bp
Locus ID: 9635
UniProt ID: Q9UQC9
Cytogenetics: 1p22.3

Domains: VWA

**Protein Families:** Druggable Genome, Ion Channels: Other, Transmembrane





### CLCA2 (NM\_006536) Human Tagged ORF Clone Lentiviral Particle - RC207731L3V

**Protein Pathways:** Olfactory transduction

MW: 103.9 kDa

Gene Summary: This gene encodes a member of the calcium-activated chloride channel regulator (CLCR)

family of proteins. Members of this family regulate the transport of chloride across the plasma membrane. The encoded protein is autoproteolytically processed to generate N- and C- terminal fragments. Expression of this gene is upregulated by the tumor suppressor protein p53 in response to DNA damage. In breast cancer, expression of this gene is

downregulated and the encoded protein may inhibit migration and invasion while promoting mesenchymal-to-epithelial transition in cancer cell lines. [provided by RefSeq, Sep 2016]