

Product datasheet for **RC207758**

Glutathione Peroxidase 3 (GPX3) (NM_002084) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Glutathione Peroxidase 3 (GPX3) (NM_002084) Human Tagged ORF Clone
Symbol: Glutathione Peroxidase 3
Synonyms: GPx-P; GSHPx-3; GSHPx-P
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC207758 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGCCCCGGCTGCTGCAGGCGTCTGCCTGCTTTCCCTGCTCCTGGCCGGCTTCGTCCTCGCAGAGCCGGG
 GACAAGAGAAGTCGAAGATGGACTGCCATGGTGGCATAAGTGGCACCATTACGAGTACGGAGCCCTCAC
 CATTGATGGGGAGGAGTACATCCCCTTCAAGCAGTATGCTGGCAAATACGTCCTTTGTCAACGTGGCC
 AGCTACTGAGGCCTGACGGGCCAGTACATTGAAGTGAATGCACACAGGAAGAGCTTGCACCATTCCGTC
 TGGTCATTCTGGGCTTTCCCTGCAACCAATTTGGAAAACAGGAACCAGGAGAGAAGTCAAGATCCCTTCC
 TACCCTCAAGTATGTCCGACCAGGTGGAGGCTTTGTCCCTAATTTCCAGCTCTTTGAGAAAAGGGGATGTC
 AATGGAGAGAAAAGAGCAGAAATTCTACACTTTCCTAAAGAAGTCTGTCTCCACCTCGGAGCTCCTGG
 GTACATCTGACCGCTCTTCTGGGAACCCATGAAGGTTACGACATCCGCTGGAAGTTTGTGAGAAAGTCT
 GGTGGGGCCAGATGGTATACCCATCATGCGCTGGCACCACCGGACCACGGTCAGCAACGTCAAGATGGAC
 ATCCTGTCCTACATGAGGCGGCAGGCAGCCCTGGGGGTCAAGAGGAAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC207758 protein sequence
 Red=Cloning site Green=Tags(s)

MARLLQASCLLSLLLAGFVSQSRGQEKSKMDCHGGISGTIYEYGALTIDGEEYIPFKQYAGKYVLFVNVA
 SY*GLTGQYIELNALQEELAPFGLVILGFPCNQFGKQEPGENSEILPTLKYYVRPGGFVVPNFQLFEKGDV
 NGEKEQKFYTLKNSCPPTSELLGTSDFLWPEPMKVHDIRWNFEKFLVGPDPGIPIMRWHHRTTVSNVKMD
 ILSYMRQAALGVKRR

TRTRPLEQKLISEEDLAANDILDYKDDDDKV



[View online »](#)

Chromatograms: https://cdn.origene.com/chromatograms/mk6267_b05.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_002084

ORF Size: 678 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#) The expression of this clone is not guaranteed due to the nature of selenoproteins.

OTI Annotation: This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.


Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_002084.5](#)

RefSeq Size:	1779 bp
RefSeq ORF:	681 bp
Locus ID:	2878
UniProt ID:	P22352
Cytogenetics:	5q33.1
Protein Families:	Druggable Genome, Secreted Protein
Protein Pathways:	Arachidonic acid metabolism, Glutathione metabolism
MW:	25.5 kDa

Gene Summary: The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyze the reduction of organic hydroperoxides and hydrogen peroxide (H₂O₂) by glutathione, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme is secreted, and is abundantly found in plasma. Downregulation of expression of this gene by promoter hypermethylation has been observed in a wide spectrum of human malignancies, including thyroid cancer, hepatocellular carcinoma and chronic myeloid leukemia. This isozyme is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2016]

Product images:



Circular map for RC207758