

Product datasheet for RC207766L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Brk (PTK6) (NM_005975) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Brk (PTK6) (NM 005975) Human Tagged ORF Clone Lentiviral Particle

Symbol: Brk
Synonyms: BRK

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_005975

 ORF Size:
 1353 bp

ORF Size: 1353 bp
ORF Nucleotide The ORF

Sequence:

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC207766).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through paturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005975.2

 RefSeq Size:
 2519 bp

 RefSeq ORF:
 1356 bp

 Locus ID:
 5753

 UniProt ID:
 013882

Cytogenetics: 20q13.33

Protein Families: Druggable Genome, Protein Kinase, Secreted Protein

MW: 51.7 kDa







Gene Summary:

The protein encoded by this gene is a cytoplasmic nonreceptor protein kinase which may function as an intracellular signal transducer in epithelial tissues. Overexpression of this gene in mammary epithelial cells leads to sensitization of the cells to epidermal growth factor and results in a partially transformed phenotype. Expression of this gene has been detected at low levels in some breast tumors but not in normal breast tissue. The encoded protein has been shown to undergo autophosphorylation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2012]