

Product datasheet for RC207775L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Parathyroid Hormone Receptor 2 (PTH2R) (NM_005048) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Parathyroid Hormone Receptor 2 (PTH2R) (NM_005048) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Parathyroid Hormone Receptor 2

Synonyms: PTHR2

Mammalian Cell Pui

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 005048

ORF Size: 1650 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207775).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 005048.2</u>

 RefSeq Size:
 2734 bp

 RefSeq ORF:
 1653 bp

 Locus ID:
 5746

 UniProt ID:
 P49190

Cytogenetics: 2q34

Domains: 7tm_2, HormR





Parathyroid Hormone Receptor 2 (PTH2R) (NM_005048) Human Tagged ORF Clone Lentiviral Particle – RC207775L3V

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

MW: 62.2 kDa

Gene Summary: The protein encoded by this gene is a member of the G-protein coupled receptor 2 family.

This protein is a receptor for parathyroid hormone (PTH). This receptor is more selective in ligand recognition and has a more specific tissue distribution compared to parathyroid hormone receptor 1 (PTHR1). It is activated only by PTH and not by parathyroid hormone-like hormone (PTHLH) and is particularly abundant in brain and pancreas. Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Jan 2013]