

Product datasheet for RC207893L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GALNT2 (NM_004481) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GALNT2 (NM_004481) Human Tagged ORF Clone Lentiviral Particle

Symbol: GALNT2

Synonyms: CDG2T; GalNAc-T2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_004481 **ORF Size:** 1713 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207893).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004481.2

 RefSeq Size:
 4484 bp

 RefSeq ORF:
 1716 bp

 Locus ID:
 2590

 UniProt ID:
 Q10471

Cytogenetics: 1q42.13

Domains: RICIN, Glycos_transf_2

Protein Families: Secreted Protein, Transmembrane





GALNT2 (NM_004481) Human Tagged ORF Clone Lentiviral Particle - RC207893L4V

Protein Pathways: Metabolic pathways, O-Glycan biosynthesis

MW: 64.7 kDa

Gene Summary: This gene encodes a member of the glycosyltransferase 2 protein family. Members of this

family initiate mucin-type O-glycoslation of peptides in the Golgi apparatus. The encoded protein may be involved in O-linked glycosylation of the immunoglobulin A1 hinge region. This gene may influence triglyceride levels, and may be involved Type 2 diabetes, as well as several types of cancer. Alternative splicing results in multiple transcript variants. [provided

by RefSeq, May 2014]