

Product datasheet for **RC207931**

YTHDC1 (NM_001031732) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	YTHDC1 (NM_001031732) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	YTHDC1
Synonyms:	YT521; YT521-B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>RC207931 representing NM_001031732
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCGGCTGACAGTCGGGAGGAGAAAGATGGAGAACTTAATGTTCTGGATGATATTTAACTGAAGTAC
 CAGAACAAGATGATGAACTGTATAATCCAGAGAGTGAACAAGATAAAAAAGGATCAAAAAG
 AAAAGTGATCGAATGGAATCTACTGATACCAAACGACAAAAGCCTTCTGTCCATTCAAGACAAGTGGTT
 TCTAAGCCACTGAGCTCATCTGTTAGCAATAACAAAAGAATAGTTAGTACAAAAGGAAAGTCAGCCACAG
 AGTATAAAAATGAGGAATATCAAAGATCTGAAAGAAAACAAGCTAGATGCTGATCGGAAAATTCGTCT
 ATCAAGTAGTGCCTCCAGAGAACCTTATAAGAATCAACCTGAAAAACCTGTGTCCGAAAAGGGATCCT
 GAAAGGAGGGCCAAATCTCTACGCCAGATGGTCTGAGAGAATTGGGCTGAAGTGGATAGACGTGCAA
 GCAGATCCAGCCAGTCTTCTAAGGAAGAAGTGAAGTCTGAAGAATATGGCTCTGACCATGAGACTGGCAG
 CAGTGGTCTTCTGATGAGCAAGGGAACAACACTGAGAATGAGGAGGAAGGAGTGAAGAAGATGTGGAG
 GAAGATGAAGAAGTAGAAGAAGATGCAGAAGAAGATGAAGAGGTGGATGAAGATGGAGAGGAGGAGGAGG
 AAGAGGAGGAGGAGGAAGAGGAGGAGGAGGAGGAGGAAGAAGAAGAAATATGAACAGGATGAGAGAGACCA
 GAAAGAGGAGGAAATGATTATGACACTCGAAGTGAAGGCACTGACTCTGGTTCTGAATCTGTTTCCCTTC
 ACAGATGGGTCTGTGAGATCTGGTTCAGGCACAGATGGATCAGATGAGAAAAAGAAGGAAAGGAGAGAG
 CTAGAGGCATATCTCAAATGTTTTGATAGAAGTGAAGCTCTGCATCAGATCATATGCAAGTTCAGAA
 AAAGAAGCATGAGAAATATCATCTTCCGTTCTGTCTGTCGAAAAGATCAAACCAAGTAACTCAAATAT
 GTGCTTCAAGATGCAAGATTTTCTCATAAAGAGTAACAACCATGAGAATGTGTCTCTTGCCAAAGCGA
 AGGGTGTATGGTCCAGCTCCCTGTAATGAGAAGAAATTAATCTTGCAATTTAGATCTGCAAGGAGTGT
 TATCTTAATATTTCTGTGAGAGAGAGTGGAAAATTTCAAGGGTTTGAAGACTTTCTTCCAGAAATCACAT
 CACGGAGGATCTCCTATACACTGGTGCTTCCAGCAGGAATGAGTGCTAAAATGCTGGGAGTGTCTTTA
 AAATTGACTGGATTTGCAGGCGTGAATTACCTTCACTAAGTCGGCTCATCTACCAATCCTTGGAAATGA
 ACATAAACCCAGTAAAGATCGGACGTGATGGACAGGAAATGAACTTGAATGTGGAACCCAGCTTTGTCTT
 CTGTTTCCCCCGATGAAAGTATTGACTTGTATCAGGTCATTCAAAAATGCGTCACAAGAGAAGAATGC
 ATTCTCAGCCCCGATCACGAGGACGTCCATCCCGTCGAGAACCAGTCCGGGATGTGGGAAGGCGTCGACC
 AGAAGATTATGATATTCATAACAGCAGAAAGAAACCAAGGATTGACTATCCCCCTGAGTTTCACCAGAGA
 CCAGGGTATTTAAAGGATCCACGATACCAGGAAGTGGACAGACGATTTTCAGGAGTTCGCCGAGATGTGT
 TTTTAAATGGGTCTACAATGATTATGTGAGGGAAATTCATAACATGGGACCACCACCCTTGGCAAGG
 AATGCCCCCTTACCCAGGAATGGAACAACCTCCACACCATCCTTACTATCAGCACCATGCTCCACCTCCT
 CAAGCTCATCCCCCTTACTCAGGACATCATCCAGTACCACATGAAGCAAGATACAGAGATAAACGAGTAC
 ATGATTATGATATGAGGGTGGATGATTTCTTCGTGCGACACAAGCTGTTGTCAGTGGCCGAGAAAGTAG
 ACCCCGTGAAAGAGACCGGGAACGAGAGCGAGACCGCCCTAGAGATAACAGACGAGACAGAGAGCGAGAT
 AGAGGACGTGATAGAGAAAGAGAAAGAGAGCGATTATGTGATCGAGACAGAGACCGAGGGGAGAGAGGTC
 GATATAGAAGA

ACGCGTACGCGGCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC207931 representing NM_001031732
 Red=Cloning site Green=Tags(s)

```

MAADSREEKDGELNVLDLITVPEQDDEL YNPESEQDKNEKKGSKRKS DRMESTDTKRQKPSVHSRQLV
SKPLSSVSNNKRI VSTKGKSATEYKNEEYQRSE RNKRLDADRKIRLSSSASREPYKNQPEKTCVRKRD
ERRAKSPTPDGSE RIGLEVDRRASRSSQSKEEVNSEEYGDHETGSSGSSDEQGNNTENEEGVEEDVE
EDEEVEEDAEEDEVEDGE EEEEEEEEEEEEEEEEEEEEEEEYE QDERDQKEEGNDYDTRSEASDSGSESVS
FDGSGVRSVSGS GTDGSEDKKERKRARGISPIVFD RSGSSASESYAGSEKKHEKLS SSVRAVRKDQTSKLY
VLQDARFFLIKSN HENVSLAKAKGVWSTLPVNEK LNLAFRSARSVILIFSVRESGKFQGFARL SESH
HGGSPIHWVLPAGMS AKMLGGVFKIDWICRREL PFKSAHLTNPWNEHKPVKIGRDGQEIELECGTQLCL
LFPPDESIDL YQVIHKMRHKRRMHSQPRSRGRPSRREPVRDVGRRRPEDYDIHNSRKKPRIDYPPEFHQR
PGYLKDPRYQEVDRRFSGVRRD VFLNGSYNDYVREFHNMGPPPPWQMPYPGMEQPPHPHPYQHHPAPP
QAHPYPYSGHHPVPEAR YRDKRVHDYDMRVDDFLRRTQAVVSGRRSRPRERDRERDRPRDNRDRERD
RGRDRERERERL CDRDRDRGERGRYRR
  
```

TRTRPLEQKLI SEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk8113_e05.zip

Restriction Sites: Sgfl-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_001031732

ORF Size: 2181 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001031732.3](#)

RefSeq Size: 6249 bp

RefSeq ORF: 2184 bp

Locus ID: 91746


UniProt ID: [Q96MU7](#)

Cytogenetics: 4q13.2

MW: 84.5 kDa

Gene Summary:

Regulator of alternative splicing that specifically recognizes and binds N6-methyladenosine (m6A)-containing RNAs (PubMed:26318451, PubMed:26876937, PubMed:25242552, PubMed:28984244). M6A is a modification present at internal sites of mRNAs and some non-coding RNAs and plays a role in the efficiency of mRNA splicing, processing and stability (PubMed:26318451, PubMed:25242552). Acts as a key regulator of exon-inclusion or exon-skipping during alternative splicing via interaction with mRNA splicing factors SRSF3 and SRSF10 (PubMed:26876937). Specifically binds m6A-containing mRNAs and promotes recruitment of SRSF3 to its mRNA-binding elements adjacent to m6A sites, leading to exon-inclusion during alternative splicing (PubMed:26876937). In contrast, interaction with SRSF3 prevents interaction with SRSF10, a splicing factor that promotes exon skipping; this prevents SRSF10 from binding to its mRNA-binding sites close to m6A-containing regions, leading to inhibit exon skipping during alternative splicing (PubMed:26876937). May also regulate alternative splice site selection (PubMed:20167602). Also involved in nuclear export of m6A-containing mRNAs via interaction with SRSF3: interaction with SRSF3 facilitates m6A-containing mRNA-binding to both SRSF3 and NXF1, promoting mRNA nuclear export (PubMed:28984244). Also recognizes and binds m6A on other RNA molecules (PubMed:27602518). Involved in random X inactivation mediated by Xist RNA: recognizes and binds m6A-containing Xist and promotes transcription repression activity of Xist (PubMed:27602518). Involved in S-adenosyl-L-methionine homeostasis by regulating expression of MAT2A transcripts, probably by binding m6A-containing MAT2A mRNAs (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for RC207931