

Product datasheet for RC208032L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ACSL3 (NM_004457) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ACSL3 (NM_004457) Human Tagged ORF Clone Lentiviral Particle

Symbol: ACSL3

Synonyms: ACS3; FACL3; LACS 3; LACS3; PRO2194

Mammalian Cell

. . .

Puromycin

Selection: Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_004457 **ORF Size:** 2160 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208032).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004457.3

 RefSeq Size:
 4369 bp

 RefSeq ORF:
 2163 bp

 Locus ID:
 2181

 UniProt ID:
 095573

Cytogenetics: 2q36.1

Domains: AMP-binding

Protein Families: Transmembrane





ACSL3 (NM_004457) Human Tagged ORF Clone Lentiviral Particle - RC208032L3V

Protein Pathways: Adipocytokine signaling pathway, Fatty acid metabolism, Metabolic pathways, PPAR signaling

pathway

MW: 80.4 kDa

Gene Summary: The protein encoded by this gene is an isozyme of the long-chain fatty-acid-coenzyme A

ligase family. Although differing in substrate specificity, subcellular localization, and tissue distribution, all isozymes of this family convert free long-chain fatty acids into fatty acyl-CoA esters, and thereby play a key role in lipid biosynthesis and fatty acid degradation. This isozyme is highly expressed in brain, and preferentially utilizes myristate, arachidonate, and eicosapentaenoate as substrates. The amino acid sequence of this isozyme is 92% identical to that of rat homolog. Two transcript variants encoding the same protein have been found for

this gene. [provided by RefSeq, Jul 2008]