

Product datasheet for RC208034L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

WASF2 (NM_006990) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: WASF2 (NM 006990) Human Tagged ORF Clone Lentiviral Particle

Symbol: WASF2

Synonyms: dJ393P12.2; IMD2; SCAR2; WASF4; WAVE2

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 006990

ORF Size: 1494 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208034).

OTI Disclaimer:

Sequence:

Domains:

er: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 006990.2

 RefSeq Size:
 4270 bp

 RefSeq ORF:
 1497 bp

 Locus ID:
 10163

 UniProt ID:
 Q9Y6W5

 Cytogenetics:
 1p36.11

Protein Families: Druggable Genome

WH2





WASF2 (NM_006990) Human Tagged ORF Clone Lentiviral Particle - RC208034L1V

Protein Pathways: Adherens junction, Fc gamma R-mediated phagocytosis, Regulation of actin cytoskeleton

MW: 54.1 kDa

Gene Summary: This gene encodes a member of the Wiskott-Aldrich syndrome protein family. The gene

product is a protein that forms a multiprotein complex that links receptor kinases and actin. Binding to actin occurs through a C-terminal verprolin homology domain in all family members. The multiprotein complex serves to tranduce signals that involve changes in cell shape, motility or function. The published map location (PMID:10381382) has been changed based on recent genomic sequence comparisons, which indicate that the expressed gene is located on chromosome 1, and a pseudogene may be located on chromosome X. Two transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jan 2011]