

Product datasheet for **RC208043L4V**

PSCD4 (CYTH4) (NM_013385) Human Tagged ORF Clone Lentiviral Particle

Product data:

| | |
|---------------------------|--|
| Product Type: | Lentiviral Particles |
| Product Name: | PSCD4 (CYTH4) (NM_013385) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | PSCD4 |
| Synonyms: | CYT4; cytohesin-4; DJ63G5.1; PSCD4 |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-mGFP-P2A-Puro (PS100093) |
| Tag: | mGFP |
| ACCN: | NM_013385 |
| ORF Size: | 1182 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC208043). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | NM_013385.2 , NP_037517.1 |
| RefSeq Size: | 3138 bp |
| RefSeq ORF: | 1185 bp |
| Locus ID: | 27128 |
| UniProt ID: | Q9UIA0 |
| Cytogenetics: | 22q13.1 |
| MW: | 45.7 kDa |


[View online »](#)

Gene Summary:

This gene encodes a member of the PSCD family of proteins, which have an N-terminal coiled-coil motif, a central Sec7 domain, and a C-terminal pleckstrin homology (PH) domain. The coiled-coil motif is involved in homodimerization, the Sec7 domain contains guanine-nucleotide exchange protein (GEP) activity, and the PH domain interacts with phospholipids and is responsible for association of PSCDs with membranes. Members of this family function as GEPs for ADP-ribosylation factors (ARFs), which are guanine nucleotide-binding proteins involved in vesicular trafficking pathways. This protein exhibits GEP activity in vitro with ARF1 and ARF5, but is inactive with ARF6. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2015]