

Product datasheet for RC208095

CD53 (NM_000560) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: CD53 (NM_000560) Human Tagged ORF Clone

Tag: Myc-DDK

Symbol: CD53

Synonyms: MOX44; TSPAN25

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC208095 representing NM_000560

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RC208095 representing NM_000560

Red=Cloning site Green=Tags(s)

MGMSSLKLLKYVLFFFNLLFWICGCCILGFGIYLLIHNNFGVLFHNLPSLTLGNVFVIVGSIIMVVAFLG CMGSIKENKCLLMSFFILLLIILLAEVTLAILLFVYEQKLNEYVAKGLTDSIHRYHSDNSTKAAWDSIQS FLQCCGINGTSDWTSGPPASCPSDRKVEGCYAKARLWFHSNFLYIGIITICVCVIEVLGMSFALTLNCQI DKTSQTIGL

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

https://cdn.origene.com/chromatograms/ja1446 c07.zip Chromatograms:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:



CTATAGGGCGGCCGC	_EcoR GGAATTCGTCGP	BamHI K		GATCTGCCG	Kozac Consensus Sgfl	C ATG -	
ORF	мми й	Miu i CG CGT ACC T R T	Not I G CGG CCG R P		Myo ag aaa cto Q K L	Tag ATC TCA I S	GAA GAG E E
GAT CTG GCA GC.		_	Flag.Tag TAC AAG G Y K	-		Fme I GTT TAA V stop	Fse I ACGGCCGGCC

^{*} The last codon before the Stop codon of the ORF

ACCN: NM 000560

ORF Size: 657 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

> of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

CD53 (NM_000560) Human Tagged ORF Clone - RC208095

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000560.4</u>

RefSeq Size: 1567 bp
RefSeq ORF: 660 bp
Locus ID: 963

 UniProt ID:
 P19397

 Cytogenetics:
 1p13.3

Domains:transmembrane4Protein Families:Transmembrane

MW: 24.2 kDa


Gene Summary: The protein encoded by this gene is a member of the transmembrane 4 superfamily, also

known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. This encoded protein is a cell surface glycoprotein that is known to complex with integrins. It contributes to the transduction of CD2-generated signals in T cells and natural killer cells and has been suggested to play a role in growth regulation. Familial deficiency of this gene has been linked to an immunodeficiency associated with recurrent infectious diseases caused by bacteria, fungi and viruses. Alternative splicing results in multiple


transcript variants. [provided by RefSeq, Mar 2016]



Product images:




Circular map for RC208095



Western blot validation of overexpression lysate (Cat# [LY424639]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208095 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).





Coomassie blue staining of purified CD53 protein (Cat# [TP308095]). The protein was produced from HEK293T cells transfected with CD53 cDNA clone (Cat# RC208095) using MegaTran 2.0 (Cat# [TT210002]).