

Product datasheet for RC208097L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

C2 (NM_000063) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: C2 (NM_000063) Human Tagged ORF Clone Lentiviral Particle

Symbol: C2

Synonyms: ARMD14; CO2

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM_000063

ORF Size: 2256 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208097).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000063.3

RefSeq Size:2862 bpRefSeq ORF:2259 bp

Locus ID: 717 **UniProt ID:** P06681

Cytogenetics: 6p21.33

Domains: CCP, Tryp_SPc, VWA

Protein Families: Druggable Genome, Protease, Secreted Protein





C2 (NM_000063) Human Tagged ORF Clone Lentiviral Particle - RC208097L3V

Protein Pathways: Complement and coagulation cascades, Systemic lupus erythematosus

MW: 83.3 kDa

Gene Summary: Component C2 is a serum glycoprotein that functions as part of the classical pathway of the

complement system. Activated C1 cleaves C2 into C2a and C2b. The serine proteinase C2a then combines with complement factor 4b to create the C3 or C5 convertase. Deficiency of C2 has been reported to associated with certain autoimmune diseases and SNPs in this gene have been associated with altered susceptibility to age-related macular degeneration. This gene localizes within the class III region of the MHC on the short arm of chromosome 6. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional transcript variants have been described in publications but their full-length

sequence has not been determined.[provided by RefSeq, Mar 2009]