

Product datasheet for RC208129L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CRKL (NM_005207) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CRKL (NM 005207) Human Tagged ORF Clone Lentiviral Particle

Symbol: CRKL

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_005207

ORF Size: 909 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208129).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 005207.2</u>

 RefSeq Size:
 5235 bp

 RefSeq ORF:
 912 bp

 Locus ID:
 1399

 UniProt ID:
 P46109

Cytogenetics: 22q11.21
Domains: SH2, SH3

Protein Families: Druggable Genome





CRKL (NM_005207) Human Tagged ORF Clone Lentiviral Particle - RC208129L4V

Protein Pathways: Chemokine signaling pathway, Chronic myeloid leukemia, ErbB signaling pathway, Fc gamma

R-mediated phagocytosis, Focal adhesion, Insulin signaling pathway, MAPK signaling pathway, Neurotrophin signaling pathway, Pathways in cancer, Regulation of actin cytoskeleton, Renal

cell carcinoma

MW: 33.6 kDa

Gene Summary: This gene encodes a protein kinase containing SH2 and SH3 (src homology) domains which

has been shown to activate the RAS and JUN kinase signaling pathways and transform fibroblasts in a RAS-dependent fashion. It is a substrate of the BCR-ABL tyrosine kinase, plays a role in fibroblast transformation by BCR-ABL, and may be oncogenic.[provided by RefSeq,

Jan 2009]