

Product datasheet for RC208170L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LRRC33 (NRROS) (NM_198565) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: LRRC33 (NRROS) (NM_198565) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: LRRC33

Synonyms: ELLP3030; GARPL1; LRRC33; SENEBAC; UNQ3030

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC208170).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_198565

ORF Size: 2076 bp



LRRC33 (NRROS) (NM_198565) Human Tagged Lenti ORF Clone - RC208170L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 198565.1, NP 940967.1

 RefSeq Size:
 2488 bp

 RefSeq ORF:
 2079 bp

 Locus ID:
 375387

 UniProt ID:
 Q86YC3

 Cytogenetics:
 3q29

Protein Families: Transmembrane

MW: 76.4 kDa

Gene Summary: Key regulator of transforming growth factor beta-1 (TGFB1) specifically required for microglia

macrophages and microglia: associates specifically via disulfide bonds with the Latency-associated peptide (LAP), which is the regulatory chain of TGFB1, and regulates integrindependent activation of TGF-beta-1 (By similarity). TGF-beta-1 activation mediated by LRRC33/NRROS is highly localized: there is little spreading of TGF-beta-1 activated from one microglial cell to neighboring microglia, suggesting the existence of localized and selective activation of TGF-beta-1 by LRRC33/NRROS (By similarity). Indirectly plays a role in Toll-like receptor (TLR) signaling: ability to inhibit TLR-mediated NF-kappa-B activation and cytokine


production is probably a consequence of its role in TGF-beta-1 signaling (PubMed:23545260).

function in the nervous system (By similarity). Required for activation of latent TGF-beta-1 in


[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for RC208170L1



Double digestion of RC208170L1 using Sgfl and Mlul