

Product datasheet for RC208199L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GALNT7 (NM_017423) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GALNT7 (NM_017423) Human Tagged ORF Clone Lentiviral Particle

Symbol: GALNT7

Synonyms: GALNAC-T7; GalNAcT7

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 017423

ORF Size: 1971 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208199).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 017423.1</u>

 RefSeq Size:
 4339 bp

 RefSeq ORF:
 1974 bp

 Locus ID:
 51809

 UniProt ID:
 Q86SF2

 Cytogenetics:
 4q34.1

Domains: RICIN, Glycos_transf_2

Protein Families: Transmembrane





GALNT7 (NM_017423) Human Tagged ORF Clone Lentiviral Particle - RC208199L3V

Protein Pathways: Metabolic pathways, O-Glycan biosynthesis

MW: 75.4 kDa

Gene Summary: This gene encodes GalNAc transferase 7, a member of the GalNAc-transferase family. The

enzyme encoded by this gene controls the initiation step of mucin-type O-linked protein glycosylation and transfer of N-acetylgalactosamine to serine and threonine amino acid residues. This enzyme is a type II transmembrane protein and shares common sequence motifs with other family members. Unlike other family members, this enzyme shows exclusive specificity for partially GalNAc-glycosylated acceptor substrates and shows no activity with non-glycosylated peptides. This protein may function as a follow-up enzyme in the initiation

step of O-glycosylation. [provided by RefSeq, Jul 2008]