

## **Product datasheet for RC208250**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Calprotectin (S100A9) (NM\_002965) Human Tagged ORF Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: Calprotectin (S100A9) (NM\_002965) Human Tagged ORF Clone

Tag: Myc-DDK Symbol: S100A9

Synonyms: 60B8AG; CAGB; CFAG; CGLB; L1AG; LIAG; MAC387; MIF; MRP14; NIF; P14

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC208250 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGACTTGCAAAATGTCGCAGCTGGAACGCAACATAGAGACCATCATCAACACCTTCCACCAATACTCTG
TGAAGCTGGGGCACCCAGACACCCTGAACCAGGGGGAATTCAAAGAGCTGGTGCGAAAAGATCTGCAAAA
TTTTCTCAAGAAGGAGAATAAGAATGAAAAGGTCATAGAACACATCATGGAGGACCTGGACACAAATGCA
GACAAGCAGCTGAGCTTCGAGGAGTTCATCATGCTGATGGCGAGGCTAACCTGGGCCTCCCACGAGAAGA
TGCACGAGGGTGACGAGGGCCCTGGCCACCACCATAAGCCAGGCCTCGGGGAGGGCACCCCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC208250 protein sequence

Red=Cloning site Green=Tags(s)

MTCKMSQLERNIETIINTFHQYSVKLGHPDTLNQGEFKELVRKDLQNFLKKENKNEKVIEHIMEDLDTNA

DKQLSFEEFIMLMARLTWASHEKMHEGDEGPGHHHKPGLGEGTP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6090">https://cdn.origene.com/chromatograms/mk6090</a> a10.zip


**Restriction Sites:** Sgfl-Mlul





#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_002965

ORF Size: 342 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

**RefSeg:** NM 002965.4

RefSeq Size: 586 bp RefSeq ORF: 345 bp



Cytogenetics:

**Locus ID:** 6280

UniProt ID: P06702

**Domains:** S\_100, EFh


1q21.3

MW: 13.2 kDa

Gene Summary: The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-

hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in the inhibition of casein kinase and altered expression of this protein is associated with the disease cystic fibrosis. This antimicrobial protein exhibits antifungal and antibacterial activity. [provided by


# **Product images:**





RefSeq, Nov 2014]

Circular map for RC208250









HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY S100A9 (Cat# RC208250, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-S100A9 (Cat# [TA804091]). Positive lysates [LY401037] (100ug) and [LC401037] (20ug) can be purchased separately from OriGene.

Western blot validation of overexpression lysate (Cat# [LY401037]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208250 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified S100A9 protein (Cat# [TP308250]). The protein was produced from HEK293T cells transfected with S100A9 cDNA clone (Cat# RC208250) using MegaTran 2.0 (Cat# [TT210002]).