

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC208304L3V

## PRAK (MAPKAPK5) (NM\_139078) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	PRAK (MAPKAPK5) (NM_139078) Human Tagged ORF Clone Lentiviral Particle
Symbol:	PRAK
Synonyms:	MAPKAP-K5; MK-5; MK5; PRAK
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_139078
ORF Size:	1656 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC208304).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 139078.1</u>
RefSeq Size:	2313 bp
RefSeq ORF:	1422 bp
Locus ID:	8550
UniProt ID:	<u>Q8IW41</u>
Cytogenetics:	12q24.12-q24.13
Domains:	pkinase
Protein Families:	Druggable Genome, Protein Kinase



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

<b>ORIGENE</b> PRAK (MAPKAPK5) (NM_139078) Human Tagged ORF Clone Lentiviral Particle – RC208304L3V	
Protein Pathways:	MAPK signaling pathway
MW:	54 kDa
Gene Summary:	The protein encoded by this gene is a tumor suppressor and member of the serine/threonine kinase family. In response to cellular stress and proinflammatory cytokines, this kinase is activated through its phosphorylation by MAP kinases including MAPK1/ERK, MAPK14/p38-alpha, and MAPK11/p38-beta. The encoded protein is found in the nucleus but translocates to the cytoplasm upon phosphorylation and activation. This kinase phosphorylates heat shock protein HSP27 at its physiologically relevant sites. Two alternately spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Nov 2012]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US