

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC208318

C19orf56 (WDR83OS) (NM_016145) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	C19orf56 (WDR83OS) (NM_016145) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	C19orf56
Synonyms:	ASTERIX; C19orf56; PTD008
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>>RC208318 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGTCCACTAACAATATGTCGGACCCACGGAGGCCGAACAAAGTGCTGAGGTACAAGCCCCCGCCGAGCG AATGTAACCCGGCCTTGGACGACCCGACGCCGGACTACATGAACCTGCTGGGCATGATCTTCAGCATGTG CGGCCTCATGCTTAAGCTGAAGTGGTGTGCTTGGGTCGCTGTCTACTGCTCCTTCATCAGCTTTGCCAAC TCTCGGAGCTCGGAGGACACGAAGCAAATGATGAGTAGCTTCATGCTGTCCATCTCTGCCGTGGTGATGT CCTATCTGCAGAATCCTCAGCCCATGACGCCCCCATGG
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG GTTTAA
Protein Sequence:	>RC208318 protein sequence <mark>Red</mark> =Cloning site Green=Tags(s)
	MSTNNMSDPRRPNKVLRYKPPPSECNPALDDPTPDYMNLLGMIFSMCGLMLKLKWCAWVAVYCSFISFAN SRSSEDTKQMMSSFMLSISAVVMSYLQNPQPMTPPW
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Chromatograms:	https://cdn.origene.com/chromatograms/mk6556_e04.zip
Restriction Sites:	Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Cloning Scheme:




* The last codon before the Stop codon of the ORF

ACCN:	NM_016145
ORF Size:	318 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 016145.4</u>
RefSeq Size:	861 bp
RefSeq ORF:	321 bp

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


	C19orf56 (WDR83OS) (NM_016145) Human Tagged ORF Clone – RC208318
Locus ID:	51398
UniProt ID:	<u>Q9Y284</u>
Cytogenetics:	19p13.13
Domains:	UPF0139
Protein Families:	Transmembrane
MW:	12.1 kDa
Gene Summary:	Component of the PAT complex, an endoplasmic reticulum (ER)-resident membrane multiprotein complex that facilitates multi-pass membrane proteins insertion into membranes (PubMed:32814900). The PAT complex acts as an intramembrane chaperone by directly interacting with nascent transmembrane domains (TMDs), releasing its substrates upon correct folding, and is needed for optimal biogenesis of multi-pass membrane proteins (PubMed:32814900). WDR83OS/Asterix is the substrate-interacting subunit of the PAT complex, whereas CCDC47 is required to maintain the stability of WDR83OS/Asterix (PubMed:12475939, PubMed:32814900). WDR83OS/Asterix associates with the first transmembrane domain (TMD1) of the nascent chain, independently of the N-glycosylation of the chain and irrespective of the amino acid sequence and transmembrane topology of TMD1 (PubMed:12475939, PubMed:32814900). The PAT complex favors the binding to TMDs with exposed hydrophilic amino acids within the lipid bilayer and provides a membrane- embedded partially hydrophilic environment in which TMD1 binds (PubMed:32814900). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RC208318

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Western blot validation of overexpression lysate (Cat# [LY414158]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208318 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US