

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC208324L4V

KIAA0427 (CTIF) (NM_014772) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	KIAA0427 (CTIF) (NM_014772) Human Tagged ORF Clone Lentiviral Particle
Symbol:	KIAA0427
Synonyms:	Gm672; KIAA0427
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_014772
ORF Size:	1794 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC208324).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 014772.2</u>
RefSeq Size:	5748 bp
RefSeq ORF:	1797 bp
Locus ID:	9811
UniProt ID:	<u>043310</u>
Cytogenetics:	18q21.1
MW:	67.6 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



KIAA0427 (CTIF) (NM_014772) Human Tagged ORF Clone Lentiviral Particle – RC208324L4V

Gene Summary:CTIF is a component of the CBP80 (NCBP1; MIM 600469)/CBP20 (NCBP2; MIM 605133)
translation initiation complex that binds cotranscriptionally to the cap end of nascent mRNA.
The CBP80/CBP20 complex is involved in a simultaneous editing and translation step that
recognizes premature termination codons (PTCs) in mRNAs and directs PTC-containing
mRNAs toward nonsense-mediated decay (NMD). On mRNAs without PTCs, the CBP80/CBP20
complex is replaced with cytoplasmic mRNA cap-binding proteins, including EIF4G (MIM
600495), and steady-state translation of the mRNAs resumes in the cytoplasm (Kim et al.,
2009 [PubMed 19648179]).[supplied by OMIM, Dec 2009]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US