

Product datasheet for **RC208350L3V**

KCNN3 (NM_170782) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	KCNN3 (NM_170782) Human Tagged ORF Clone Lentiviral Particle
Symbol:	KCNN3
Synonyms:	hSK3; KCa2.3; SK3; SKCA3; ZLS3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_170782
ORF Size:	1278 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC208350).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_170782.1
RefSeq Size:	11956 bp
RefSeq ORF:	1281 bp
Locus ID:	3782
UniProt ID:	Q9UGI6
Cytogenetics:	1q21.3
Protein Families:	Druggable Genome, Ion Channels: Potassium, Transmembrane
MW:	48.1 kDa



[View online »](#)

Gene Summary:

Action potentials in vertebrate neurons are followed by an afterhyperpolarization (AHP) that may persist for several seconds and may have profound consequences for the firing pattern of the neuron. Each component of the AHP is kinetically distinct and is mediated by different calcium-activated potassium channels. This gene belongs to the KCNN family of potassium channels. It encodes an integral membrane protein that forms a voltage-independent calcium-activated channel, which is thought to regulate neuronal excitability by contributing to the slow component of synaptic AHP. This gene contains two CAG repeat regions in the coding sequence. It was thought that expansion of one or both of these repeats could lead to an increased susceptibility to schizophrenia or bipolar disorder, but studies indicate that this is probably not the case. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2011]