

Product datasheet for RC208404L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SREBP1 (SREBF1) (NM 004176) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SREBP1 (SREBF1) (NM_004176) Human Tagged ORF Clone Lentiviral Particle

Symbol:

bHLHd1; HMD; IFAP2; SREBP1 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

mGFP Tag:

NM 004176 ACCN: **ORF Size:** 3441 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208404).

Sequence: OTI Disclaimer:

Domains:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 004176.3

RefSeq Size: 4922 bp RefSeq ORF: 3444 bp Locus ID: 6720 **UniProt ID:** P36956 Cytogenetics: 17p11.2

HLH **Protein Families:** Druggable Genome, Transcription Factors





Protein Pathways: Insulin signaling pathway

MW: 122.2 kDa

Gene Summary: This gene encodes a basic helix-loop-helix-leucine zipper (bHLH-Zip) transcription factor that

binds to the sterol regulatory element-1 (SRE1), which is a motif that is found in the promoter of the low density lipoprotein receptor gene and other genes involved in sterol biosynthesis. The encoded protein is synthesized as a precursor that is initially attached to the nuclear membrane and endoplasmic reticulum. Following cleavage, the mature protein translocates to the nucleus and activates transcription. This cleaveage is inhibited by sterols. This gene is located within the Smith-Magenis syndrome region on chromosome 17. Alternative promoter usage and splicing result in multiple transcript variants, including SREBP-1a and SREBP-1c, which correspond to RefSeq transcript variants 2 and 3, respectively. [provided by RefSeq,

Nov 2017]