

Product datasheet for RC208561L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

WNT10A (NM_025216) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: WNT10A (NM_025216) Human Tagged ORF Clone Lentiviral Particle

Symbol: WNT10A

Synonyms: OODD; SSPS; STHAG4

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_025216

ORF Size: 1251 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208561).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 025216.2

 RefSeq Size:
 2375 bp

 RefSeq ORF:
 1254 bp

 Locus ID:
 80326

 UniProt ID:
 Q9GZT5

 Cytogenetics:
 2q35

Protein Families: Secreted Protein, Transmembrane





WNT10A (NM_025216) Human Tagged ORF Clone Lentiviral Particle - RC208561L1V

Protein Pathways: Basal cell carcinoma, Hedgehog signaling pathway, Melanogenesis, Pathways in cancer, Wnt

signaling pathway

MW: 42.4 kDa

Gene Summary: The WNT gene family consists of structurally related genes which encode secreted signaling

proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It is strongly expressed in the cell lines of promyelocytic leukemia and Burkitt's lymphoma. In addition, it and another family member, the WNT6 gene, are strongly coexpressed in colorectal cancer cell lines. The gene overexpression may play key roles in carcinogenesis through activation of the WNT-beta-catenin-TCF signaling pathway. This gene and the WNT6 gene are clustered in the chromosome 2q35 region.

[provided by RefSeq, Jul 2008]