

## Product datasheet for **RC208561L3V**

### WNT10A (NM\_025216) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	WNT10A (NM_025216) Human Tagged ORF Clone Lentiviral Particle
Symbol:	WNT10A
Synonyms:	OODD; SSPS; STHAG4
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_025216
ORF Size:	1251 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC208561).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_025216.2</a>
RefSeq Size:	2375 bp
RefSeq ORF:	1254 bp
Locus ID:	80326
UniProt ID:	<a href="#">Q9GZT5</a>
Cytogenetics:	2q35
Protein Families:	Secreted Protein, Transmembrane



[View online »](#)

<b>Protein Pathways:</b>	Basal cell carcinoma, Hedgehog signaling pathway, Melanogenesis, Pathways in cancer, Wnt signaling pathway
<b>MW:</b>	42.4 kDa
<b>Gene Summary:</b>	<p>The WNT gene family consists of structurally related genes which encode secreted signaling proteins. These proteins have been implicated in oncogenesis and in several developmental processes, including regulation of cell fate and patterning during embryogenesis. This gene is a member of the WNT gene family. It is strongly expressed in the cell lines of promyelocytic leukemia and Burkitt's lymphoma. In addition, it and another family member, the WNT6 gene, are strongly coexpressed in colorectal cancer cell lines. The gene overexpression may play key roles in carcinogenesis through activation of the WNT-beta-catenin-TCF signaling pathway. This gene and the WNT6 gene are clustered in the chromosome 2q35 region. [provided by RefSeq, Jul 2008]</p>