

Product datasheet for RC208605L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CCR1 (NM_001295) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CCR1 (NM_001295) Human Tagged ORF Clone Lentiviral Particle

Symbol: CCR²

Synonyms: CD191; CKR-1; CKR1; CMKBR1; HM145; MIP1aR; SCYAR1

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 001295

ORF Size: 1065 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC208605).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001295.2</u>

 RefSeq Size:
 2690 bp

 RefSeq ORF:
 1068 bp

 Locus ID:
 1230

 UniProt ID:
 P32246

 Cytogenetics:
 3p21.31

Domains: 7tm_1

Protein Families: Druggable Genome, GPCR, Transmembrane





CCR1 (NM_001295) Human Tagged ORF Clone Lentiviral Particle - RC208605L1V

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction

MW: 41.2 kDa

Gene Summary: This gene encodes a member of the beta chemokine receptor family, which is predicted to be

a seven transmembrane protein similar to G protein-coupled receptors. The ligands of this receptor include macrophage inflammatory protein 1 alpha (MIP-1 alpha), regulated on activation normal T expressed and secreted protein (RANTES), monocyte chemoattractant protein 3 (MCP-3), and myeloid progenitor inhibitory factor-1 (MPIF-1). Chemokines and their receptors mediated signal transduction are critical for the recruitment of effector immune cells to the site of inflammation. Knockout studies of the mouse homolog suggested the roles of this gene in host protection from inflammatory response, and susceptibility to virus and parasite. This gene and other chemokine receptor genes, including CCR2, CCRL2, CCR3, CCR5 and CCXCR1, are found to form a gene cluster on chromosome 3p. [provided by RefSeq, Jul

2008]