

Product datasheet for RC208634L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

EXOSC6 (NM_058219) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: EXOSC6 (NM_058219) Human Tagged ORF Clone Lentiviral Particle

Symbol: EXOSC6

Synonyms: EAP4; hMtr3p; MTR3; Mtr3p; p11

Mammalian Cell

Selection:

ACCN:

None

NM 058219

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ORF Size: 816 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC208634).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 058219.2</u>

 RefSeq Size:
 1729 bp

 RefSeq ORF:
 819 bp

 Locus ID:
 118460

 UniProt ID:
 Q5RKV6

Cytogenetics: 16q22.1

Domains: RNase_PH_C

Protein Pathways: RNA degradation





ORIGENE

MW: 28.2 kDa

Gene Summary: This ge

This gene product constitutes one of the subunits of the multisubunit particle called exosome, which mediates mRNA degradation. The composition of human exosome is similar to its yeast counterpart. This protein is homologous to the yeast Mtr3 protein. Its exact function is not known, however, it has been shown using a cell-free RNA decay system that the exosome is required for rapid degradation of unstable mRNAs containing AU-rich elements (AREs), but not for poly(A) shortening. The exosome does not recognize ARE-containing mRNAs on its own, but requires ARE-binding proteins that could interact with the exosome and recruit it to unstable mRNAs, thereby promoting their rapid degradation. [provided by RefSeq, Jul 2008]