

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC208655

## H3FD (HIST1H3E) (NM\_003532) Human Tagged ORF Clone


## **Product data:**

Product Type:	Expression Plasmids
Product Name:	H3FD (HIST1H3E) (NM_003532) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	H3FD
Synonyms:	H3.1; H3/d; H3C1; H3C2; H3C3; H3C4; H3C7; H3C8; H3C10; H3C11; H3C12; H3FD; HIST1H3E
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>&gt;RC208655 representing NM_003532 Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCGCGTACTAAGCAGACGGCTCGTAAATCCACAGGCGGTAAAGCACCGCGCAAAACAGCTGGCCACTA AGGCAGCTCGCAAGAGCGCTCCGGCCACGGGCGGCGTGAAGAAGCCCCATCGCTACCGCCCTGGCACCGT GGCTCTGCGCGAGATCCGTCGCTACCAGAAGTCTACCGAGCTTCTAATCCGGAAGCTGCCGTTTCAGCGC CTGGTGCGAGAAATAGCTCAGGACTTCAAGACCGACCTGCGCTTCCAGAGTTCCGCGGTGATGGCGCTGC AGGAGGCCTGCGAGGCCTACTTGGTGGGGGCTTTTCGAGGACACCAACCTGTGCGCTATTCATGCCAAACG CGTGACCATCATGCCTAAAGACATCCAGCTTGCCGCCGCCGCATTCGTGGGGAGAGGGCG
	AGCGGACCGACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC TGGATTACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	<pre>&gt;RC208655 representing NM_003532 Red=Cloning site Green=Tags(s)</pre>
	MARTKQTARKSTGGKAPRKQLATKAARKSAPATGGVKKPHRYRPGTVALREIRRYQKSTELLIRKLPFQR LVREIAQDFKTDLRFQSSAVMALQEACEAYLVGLFEDTNLCAIHAKRVTIMPKDIQLARRIRGERA
	SGPTRTRRLEQKLISEEDLAANDILDYKDDDDKV
Chromatograms:	https://cdn.origene.com/chromatograms/mk6616_b04.zip
Restriction Sites:	Sgfl-RsrII



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **Cloning Scheme:**




ACCN:	NM_003532
ORF Size:	408 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 003532.3</u>
RefSeq Size:	462 bp
RefSeq ORF:	411 bp
Locus ID:	8353
	<u>P68431</u>


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Service March (HIST1H3E) (NM_003532) Human Tagged ORF Clone – RC208655		
Cytogenetics:	6p22.2	
Domains:	H3, histone	
Protein Pathways:	Systemic lupus erythematosus	
MW:	15.2 kDa	
Gene Summary:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6. [provided by RefSeq, Aug 2015]	

## **Product images:**



Circular map for RC208655



Western blot validation of overexpression lysate (Cat# [LY418620]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208655 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US