

Product datasheet for RC208663L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LEF1 (NM_016269) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: LEF1 (NM_016269) Human Tagged ORF Clone Lentiviral Particle

Symbol: LEF

Synonyms: LEF-1; TCF1ALPHA; TCF7L3; TCF10

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM_016269

ORF Size: 1197 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208663).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 016269.2

 RefSeq Size:
 3620 bp

 RefSeq ORF:
 1200 bp

 Locus ID:
 51176

 UniProt ID:
 Q9UJU2

 Cytogenetics:
 4q25

 Domains:
 HMG

Protein Families: Adult stem cells, Druggable Genome, ES Cell Differentiation/IPS, Transcription Factors





LEF1 (NM_016269) Human Tagged ORF Clone Lentiviral Particle - RC208663L1V

Protein Pathways: Acute myeloid leukemia, Adherens junction, Arrhythmogenic right ventricular

cardiomyopathy (ARVC), Basal cell carcinoma, Colorectal cancer, Endometrial cancer,

Melanogenesis, Pathways in cancer, Prostate cancer, Thyroid cancer, Wnt signaling pathway

MW: 44.2 kDa

Gene Summary: This gene encodes a transcription factor belonging to a family of proteins that share

homology with the high mobility group protein-1. The protein encoded by this gene can bind to a functionally important site in the T-cell receptor-alpha enhancer, thereby conferring maximal enhancer activity. This transcription factor is involved in the Wnt signaling pathway, and it may function in hair cell differentiation and follicle morphogenesis. Mutations in this gene have been found in somatic sebaceous tumors. This gene has also been linked to other cancers, including androgen-independent prostate cancer. Alternative splicing results in

multiple transcript variants. [provided by RefSeq, Oct 2009]