

Product datasheet for RC208673L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

BMPR2 (NM_001204) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: BMPR2 (NM 001204) Human Tagged ORF Clone Lentiviral Particle

Symbol: BMPR2

Synonyms: BMPR-II; BMPR3; BMR2; BRK-3; POVD1; PPH1; T-ALK

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001204 **ORF Size:** 3114 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208673).

OTI Disclaimer:

Cytogenetics:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001204.5</u>

RefSeq Size:12086 bpRefSeq ORF:3117 bp

Locus ID: 659 **UniProt ID:** Q13873

Domains: Activin_recp, pkinase, TyrKc, S_TKc

2q33.1-q33.2

Protein Families: Druggable Genome, Protein Kinase, Transmembrane





BMPR2 (NM_001204) Human Tagged ORF Clone Lentiviral Particle - RC208673L4V

Protein Pathways: Cytokine-cytokine receptor interaction, TGF-beta signaling pathway

MW: 115.2 kDa

Gene Summary: This gene encodes a member of the bone morphogenetic protein (BMP) receptor family of

transmembrane serine/threonine kinases. The ligands of this receptor are members of the

TGF-beta superfamily. BMPs are involved in endochondral bone formation and

embryogenesis. These proteins transduce their signals through the formation of heteromeric complexes of two different types of serine (threonine) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Mutations in this gene have been associated with primary pulmonary hypertension, both familial and fenfluramine-associated,

and with pulmonary venoocclusive disease. [provided by RefSeq, May 2020]