

Product datasheet for RC208741L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

UBE2C (NM_007019) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: UBE2C (NM_007019) Human Tagged ORF Clone Lentiviral Particle

Symbol: UBE2C

Synonyms: dJ447F3.2; UBCH10

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_007019

ORF Size: 537 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208741).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 007019.2</u>

 RefSeq Size:
 864 bp

 RefSeq ORF:
 540 bp

 Locus ID:
 11065

 UniProt ID:
 000762

 Cytogenetics:
 20q13.12

 Domains:
 UBCc

Protein Families: Druggable Genome





UBE2C (NM_007019) Human Tagged ORF Clone Lentiviral Particle - RC208741L2V

Protein Pathways: Ubiquitin mediated proteolysis

MW: 19.7 kDa

Gene Summary: The modification of proteins with ubiquitin is an important cellular mechanism for targeting

abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013]