

Product datasheet for RC208761L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RbAp48 (RBBP4) (NM 005610) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RbAp48 (RBBP4) (NM_005610) Human Tagged ORF Clone Lentiviral Particle

Symbol: RbAp48

Synonyms: lin-53; NURF55; RBAP48

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_005610

 ORF Size:
 1275 bp

ORF Nucleotide

Sequence:

Domains:

The ORF insert of this clone is exactly the same as(RC208761).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 005610.1, NP 005601.1

 RefSeq Size:
 7943 bp

 RefSeq ORF:
 1278 bp

 Locus ID:
 5928

 UniProt ID:
 Q09028

 Cytogenetics:
 1p35.1

Protein Families: Druggable Genome, Transcription Factors

WD40





ORIGENE

MW: 47.7 kDa

Gene Summary:

This gene encodes a ubiquitously expressed nuclear protein which belongs to a highly conserved subfamily of WD-repeat proteins. It is present in protein complexes involved in histone acetylation and chromatin assembly. It is part of the Mi-2 complex which has been implicated in chromatin remodeling and transcriptional repression associated with histone deacetylation. This encoded protein is also part of co-repressor complexes, which is an integral component of transcriptional silencing. It is found among several cellular proteins that bind directly to retinoblastoma protein to regulate cell proliferation. This protein also seems to be involved in transcriptional repression of E2F-responsive genes. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2008]