

Product datasheet for RC208769L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DDX1 (NM_004939) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: DDX1 (NM 004939) Human Tagged ORF Clone Lentiviral Particle

Symbol: DDX'

Synonyms: DBP-RB; UKVH5d

Mammalian Cell

Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_004939 **ORF Size:** 2220 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208769).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 004939.1

 RefSeq Size:
 2755 bp

 RefSeq ORF:
 2223 bp

 Locus ID:
 1653

 UniProt ID:
 Q92499

 Cytogenetics:
 2p24.3

Domains: DEAD, helicase_C, SPRY

Protein Families: Druggable Genome





ORIGENE

MW: 82.4 kDa

Gene Summary: DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are

putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and

mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis,

spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein of unknown function. It shows high transcription levels in 2 retinoblastoma cell lines and in

tissues of neuroectodermal origin. [provided by RefSeq, Jul 2008]