

Product datasheet for **RC208776**

MID1 (NM_000381) Human Tagged ORF Clone

Product data:

| | |
|---------------------------|---|
| Product Type: | Expression Plasmids |
| Product Name: | MID1 (NM_000381) Human Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | MID1 |
| Synonyms: | BBBG1; FXY; GBBB1; MIDIN; OGS1; OS; OSX; RNF59; TRIM18; XPRF; ZNFXY |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |



[View online »](#)

ORF Nucleotide
Sequence:

>RC208776 ORF sequence
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGGAACACTGGAGTCAGAAGTACCTGACCTGCCCATTGTTGTCTGGAGCTCTTTGAGGACCTCTTCTACTGC
CCTGCGCACACAGCCTCTGCTTCAACTGCGCCACCGCATCCTAGTATCACACTGTGCCACCAACGAGTC
TGTGGAGTCCATCACCGCCTTCCAGTGCACCCACCTGCCGGCATGTCATCACCCCTCAGCCAGCGAGGTCTA
GACGGGCTCAAGCGCAACGTCACCCACAGAACATCATCGACAGGTTCCAGAAAGCATCAGTGAGCGGGC
CCAACCTCTCCAGCGAGACCCGTCGGGAGCGGGCCTTTGACGCCAACACCATGACCTCCGCCGAGAAGGT
CCTCTGCCAGTTTTGTGACCAGGATCCTGCCAGGACGCTGTGAAGACCTGTGTCACCTGTGAAGTATCC
TACTGTGACGAGTGCCTGAAAGCCACTACCCGAATAAGAAGCCCTTTACAGGCCATCGTCTGATTGAGC
CAATCCGGACTCTCACATCCGGGGCTGATGTGCTTGGAGCATGAGGATGAGAAGTGAATATGTACTG
TGTGACCGATGACCAGTTAATCTGTGCCTTGTGTAAGTGGTGGGCGGCACCGCATCATCAGGTGGCA
GCTTTGAGTGAGCGCTATGACAAATTGAAGCAAACCTTAGAGAGTAACCTACCAACCTTATTAAGAGGA
ACACAGAACTGGAGACCTTTTGGCTAAACTCATCAAACCTGTCAACATGTTGAAGTCAATGCATCACG
TCAAGAAGCCAAATTGACAGAGGAGTGTGATCTTCTCATTGAGATCATTAGCAAAAGACGACAGATTATT
GGAACCAAGATCAAAGAAGGGAAGGTGATGAGGCTTCGAAAACCTGGCTCAGCAGATTGCAAACTGCAAAAC
AGTGCAATTGAGCGGTGAGCATCACTCATCTCCAAAGCGGAACTCTCTGAAGGAGAATGATCATGCGCG
TTTCTACAGACTGCTAAGAATATCACCGAGAGTCTCCATGGCAACTGCATCCTCCAGGTTCTAATT
CCTGAAATCAACCTCAATGACACATTTGACACCTTTGCCTTAGATTTTTCCCGAGAGAAGAACTGCTAG
AATGCTGGATTACCTTACAGCTCCCAACCTCCACAATTAGAGAAGAGCTCTGCACAGCTTCATATGA
CACCATCACTGTGATTGGACCTCCGATGATGAGTTCAGCGTGGTCTCCTACGAGCTCCAGTACACCATA
TTCACCGACAAGCCAACGTCGTTAGTCTGTGTAATTCGGCTGATAGCTGGATGATAGTACCCAACATCA
AGCAGAACCCTACACGGTGCACGGTCTGCAGAGCGGCACCAAGTACATCTTCATGGTCAAGGCCATCAA
CCAGGGCGGCAGCCGAGCAGTGAAGCTGGGAAGTTGAAGACAAACAGCCAACCATTTAAACTGGATCCC
AAATCTGCTCATCGAAAACCTGAAGGTGTCCATGATAACTTGACAGTAGAACGTGATGAGTCATCATCCA
AGAAGAGTCACACACCTGAACGCTTCACCAGCCAGGGGAGCTATGGAGTAGCTGGAAATGTGTTTATTGA
TAGTGGCCGGCATTATTGGGAAGTGGTCATAAGTGAAGCACATGGTATGCCATTGGTCTTGCTTACAAA
TCAGCCCCGAAGCATGAATGGATTGGGAAGAACTCTGCTTCTGGGCGCTCTGCCGCTGCAACAATAACT
GGGTGGTGAAGACAATAGCAAGGAAATCCCCATTGAGCCTGCCCCACCTCCGGCGCGTGGGCATCCT
GCTGGACTATGATAACGGCTCTATCGCCTTTTATGATGCTTTGAACTCCATCCACCTCTACACCTTCGAC
GTCGCATTTGCGCAGCCTGTTTGGCCACCTTCACCGTGTGGAACAAGTGTCTGACGATTATCACTGGGC
TCCCTATCCCAGACCATTGGACTGCACAGAGCAGCTGCCG

ACGCGTACGCGGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC208776 protein sequence
Red=Cloning site Green=Tags(s)

```
METLESELTCPICLELFEDPLLLPCAHSLCFNCAHRILVSHCATNESVESITAFQCPTCRHVITLSQRGL
DGLKRNVTLQNIIDRFQKASVSGPNPSETRRERAFDANTMTSAEKVLCQFCDQDPAQDAVKTCVTCVSV
YCDECLKATHPNKKPFTGHRLIEPIPDSHIRGLMCLHEHEDEKVNMYCVTDDQLICALCKLVGRHRDHQVA
ALSERYDKLKQNLNLESNLTNLIKRNTELETLLAKLIQTQCQHVEVNASRQEAKLTEECDLLIEIIQRRQII
GTKIKEGKVMRLRKLAAQIQIANCKQCIERSASLISQAEHSLKENDHARFLQTAKNITERVSMATASSQVLI
PEINLNDTFDFALDFSREKLLLECLDYLTAPNPPTIREELCTASYDTITVHWTSDDEFVSVSYELQYTI
FTGQANVVSLSADSWMIVPNIKQNHVTVHGLQSGTKYIFMVKAINQAGRSRSEPGKLTNSQPFLKDP
KSAHRKLVSHDNLTVRDESSSKSHTPERFTSQGSYGVAGNVFIDSGRHYWEVVISGSTWYAIGLAYK
SAPKHEWIGKNSASWALCRCNNWVVRHNSKEIPIEPAPHLRRVGILLDYDNGSIAFYDALNSIHLTYTFD
VAFAQPVCPTFTVWNKCLTIITGLPIPDLHDCTEQLP
```

TRTRPLEQKLISEEDLANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6218_c06.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_000381

ORF Size: 2001 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_000381.4](#)

RefSeq Size: 6463 bp

RefSeq ORF: 2004 bp

Locus ID: 4281

UniProt ID: [O15344](#)

Cytogenetics: Xp22.2

Domains: zf-B_box, RING, BBC, SPRY, FN3


Protein Families: Druggable Genome

Protein Pathways: Ubiquitin mediated proteolysis


MW: 75.3 kDa

Gene Summary: The protein encoded by this gene is a member of the tripartite motif (TRIM) family, also known as the 'RING-B box-coiled coil' (RBCC) subgroup of RING finger proteins. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This protein forms homodimers which associate with microtubules in the cytoplasm. The protein is likely involved in the formation of multiprotein structures acting as anchor points to microtubules. Mutations in this gene have been associated with the X-linked form of Opitz syndrome, which is characterized by midline abnormalities such as cleft lip, laryngeal cleft, heart defects, hypospadias, and agenesis of the corpus callosum. This gene was also the first example of a gene subject to X inactivation in human while escaping it in mouse. Alternative promoter use, alternative splicing and alternative polyadenylation result in multiple transcript variants that have different tissue specificities. [provided by RefSeq, Dec 2016]


Product images:



Circular map for RC208776



Western blot validation of overexpression lysate (Cat# [LY424750]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208776 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified MID1 protein (Cat# [TP308776]). The protein was produced from HEK293T cells transfected with MID1 cDNA clone (Cat# RC208776) using MegaTran 2.0 (Cat# [TT210002]).