

Product datasheet for RC208789L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CKII alpha (CSNK2A1) (NM_177560) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CKII alpha (CSNK2A1) (NM_177560) Human Tagged ORF Clone Lentiviral Particle

Symbol: CSNK2A1

Synonyms: CK2A1; Cka1; Cka2; CKII; OCNDS

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_177560 **ORF Size:** 1173 bp

ORF Nucleotide

. . , 5 5 5

Sequence:

Cytogenetics:

The ORF insert of this clone is exactly the same as(RC208789).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 177560.2, NP 808228.1

20p13

 RefSeq Size:
 2522 bp

 RefSeq ORF:
 768 bp

 Locus ID:
 1457

 UniProt ID:
 P68400

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Protein Kinase

Protein Pathways: Adherens junction, Tight junction, Wnt signaling pathway





MW: 45.1 kDa

Gene Summary:

Casein kinase II is a serine/threonine protein kinase that phosphorylates acidic proteins such as casein. It is involved in various cellular processes, including cell cycle control, apoptosis, and circadian rhythm. The kinase exists as a tetramer and is composed of an alpha, an alphaprime, and two beta subunits. The alpha subunits contain the catalytic activity while the beta subunits undergo autophosphorylation. The protein encoded by this gene represents the alpha subunit. Multiple transcript variants encoding different protein isoforms have been found for this gene. [provided by RefSeq, Apr 2018]