

Product datasheet for RC208816L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ZNF384 (NM_001039920) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ZNF384 (NM_001039920) Human Tagged ORF Clone Lentiviral Particle

Symbol: ZNF384

Synonyms: CAGH1; CAGH1A; CIZ; ERDA2; NMP4; NP; TNRC1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001039920

ORF Size: 1347 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC208816).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 001039920.1

 RefSeq Size:
 2903 bp

 RefSeq ORF:
 1386 bp

 Locus ID:
 171017

 UniProt ID:
 Q8TF68

 Cytogenetics:
 12p13.31

Protein Families: Transcription Factors

MW: 50.4 kDa







Gene Summary:

This gene encodes a C2H2-type zinc finger protein, which may function as a transcription factor. This gene also contains long CAG trinucleotide repeats that encode consecutive glutamine residues. The protein appears to bind and regulate the promoters of the extracellular matrix genes MMP1, MMP3, MMP7 and COL1A1. Studies in mouse suggest that nuclear matrix transcription factors (NP/NMP4) may be part of a general mechanical pathway that couples cell construction and function during extracellular matrix remodeling. Alternative splicing results in multiple transcript variants. Recurrent rearrangements of this gene with the Ewing's sarcoma gene, EWSR1 on chromosome 22, or with the TAF15 gene on chromosome 17, or with the TCF3 (E2A) gene on chromosome 19, have been observed in acute leukemia. A related pseudogene has been identified on chromosome 7. [provided by RefSeq, Apr 2011]