

Product datasheet for RC208857

HBQ1 (NM_005331) Human Tagged ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	HBQ1 (NM_005331) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	HBQ1
Synonyms:	HBQ
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	<pre>>RC208857 ORF sequence Red=Cloning site Blue=ORF Green=Tags(s)</pre>
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCGCTGTCCGCGGAGGACCGGGCGCTGGTGCGCGCCCTGTGGAAGAAGCTGGGCAGCAACGTCGGCG TCTACACGACAGAGGCCCTGGAAAGGACCTTCCTGGCTTTCCCCGCCACGAAGACCTACTTCTCCCACCT GGACCTGAGCCCCGGCTCCTCACAAGTCAGAGCCCACGGCCAGGAAGGTGGCGGACGCGCTGAGCCTCGCC GTGGAGCGCCTGGACGACCTACCCCACGCGCTGTCCGCGCTGAGCCACCTGCACGCGTGCCAGCTGCGAG TGGACCCGGCCAGCTTCCAGCTCCTGGGCCACTGCCTGCTGGTAACCCTCGCCGGCACTACCCCGGAGA CTTCAGCCCCGCGCTGCAGGCGTCGCTGGACAAGTTCCTGAGCCACGTTATCTCGGCGCTGGTTTCCGAG TACCGC
	ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGA TTACAAGGATGACGACGATAAGGTTTAA
Protein Sequence:	<pre>>RC208857 protein sequence Red=Cloning site Green=Tags(s)</pre>
	MALSAEDRALVRALWKKLGSNVGVYTTEALERTFLAFPATKTYFSHLDLSPGSSQVRAHGQKVADALSLA VERLDDLPHALSALSHLHACQLRVDPASFQLLGHCLLVTLARHYPGDFSPALQASLDKFLSHVISALVSE YR
	TRPLEQKLISEEDLAANDILDYKDDDDKV
Chromatograms:	https://cdn.origene.com/chromatograms/mk6798_c08.zip




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	1 (NM_005331) Human Tagged ORF Clone – RC208857
Restriction Sites:	Sgfl-Notl
Cloning Scheme:	Cloning sites used for ORF Shuttling: Sgf1 ORF Not I GCGATCGC C ATG 18819 AC G CGG CCG C TC Kozac
	<u>DL</u> AAN DIL <u>DYK DDDK</u> V stoop.
ACCN:	NM_005331
ORF Size:	426 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Metho	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 005331.5</u>
RefSeq Size:	653 bp
RefSeq ORF:	429 bp


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	HBQ1 (NM_005331) Human Tagged ORF Clone – RC208857
Locus ID:	3049
UniProt ID:	<u>P09105</u>
Cytogenetics:	16p13.3
MW:	15.5 kDa
Gene Summary:	Theta-globin mRNA is found in human fetal erythroid tissue but not in adult erythroid or other nonerythroid tissue. The theta-1 gene may be expressed very early in embryonic life, perhaps sometime before 5 weeks. Theta-1 is a member of the human alpha-globin gene cluster that involves five functional genes and two pseudogenes. The order of genes is: 5' - zeta - pseudozeta - mu - pseudoalpha-2 -pseudoalpha-1 - alpha-2 - alpha-1 - theta-1 - 3'. Research supports a transcriptionally active role for the gene and a functional role for the peptide in specific cells, possibly those of early erythroid tissue. [provided by RefSeq, Jul 2008]

Product images:




Circular map for RC208857



Western blot validation of overexpression lysate (Cat# [LY417378]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208857 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Coomassie blue staining of purified HBQ1 protein (Cat# [TP308857]). The protein was produced from HEK293T cells transfected with HBQ1 cDNA clone (Cat# RC208857) using MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US