

## **Product datasheet for RC208874**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Angiogenin (ANG) (NM\_001145) Human Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Angiogenin (ANG) (NM\_001145) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: Angiogenin

Synonyms: ALS9; HEL168; RAA1; RNASE4; RNASE5

Mammalian Cell N

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC208874 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGTGATGGCCTGGGCGTTTTGTTGTTGTTCTTCGTGCTGGGTCTGGGTCTGACCCCACCGACCCTGG CTCAGGATAACTCCAGGTACACACACTTCCTGACCCAGCACTATGATGCCAAACCACAGGGCCGGGATGA CAGATACTGTGAAAGCATCATGAGGAGACGGGGCCTGACCTCACCCTGCAAAGACATCAACACATTTATT CATGGCAACAAGCGCAGCATCAAGGCCATCTGTGAAAACAAGAATGGAAACCCTCACAGAGAAAACCTAA GAATAAGCAAGTCTTCTTTCCAGGTCACCACTTGCAAGCTACATGGAGGTTCCCCCTGGCCTCCATGCCA GTACCGAGCCACAGCGGGGTTCAGAAACGTTGTTGTTGCTTGTAAAAATGGCTTACCTGTCCACTTGGAT CAGTCAATTTTCCGTCGTCCG

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC208874 protein sequence

Red=Cloning site Green=Tags(s)

MVMGLGVLLLVFVLGLGLTPPTLAQDNSRYTHFLTQHYDAKPQGRDDRYCESIMRRRGLTSPCKDINTFI HGNKRSIKAICENKNGNPHRENLRISKSSFQVTTCKLHGGSPWPPCQYRATAGFRNVVVACENGLPVHLD

**QSIFRRP** 

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Chromatograms:** https://cdn.origene.com/chromatograms/mk6035\_e08.zip






**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_001145

ORF Size: 441 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 001145.4</u>

RefSeq Size: 1222 bp RefSeq ORF: 444 bp



Locus ID: 283

UniProt ID: P03950

Cytogenetics: 14q11.2

Domains: RNAse\_Pc


**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**MW:** 16.6 kDa

**Gene Summary:** The protein encoded by this gene is a member of the RNase A superfamily though it has


relatively weak ribonucleolytic activity. This protein is a potent mediator of new blood vessel formation and thus, in addition to the name RNase5, is commonly called angiogenin. This protein induces angiogenesis after binding to actin on the surface of endothelial cells. This protein also accumulates at the nucleolus where it stimulates ribosomal transcription. Under stress conditions this protein translocates to the cytosol where it hydrolyzes cellular tRNAs and influences protein synthesis. A signal peptide is cleaved from the precursor protein to produce a mature protein which contains a nuclear localization signal, a cell binding motif, and a catalytic domain. This protein has been shown to be both neurotrophic and neuroprotective and the mature protein has antimicrobial activity against some bacteria and fungi, including S. pneumoniae and C. albicans. Due to its effect on rRNA production and angiogenesis this gene plays important roles in cell growth and tumor progression. Mutations in this gene are associated with progression of amyotrophic lateral sclerosis (ALS). This gene and the neighboring RNase4 gene share promoters and 5' exons though each gene then splices to a distinct 3' exon containing the complete coding region of each gene. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2020]


## **Product images:**



Circular map for RC208874







Western blot validation of overexpression lysate (Cat# [LY420124]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with [RC223486] using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified ANG protein (Cat# [TP308874]). The protein was produced from HEK293T cells transfected with ANG cDNA clone (Cat# RC208874) using MegaTran 2.0 (Cat# [TT210002]).