

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RC208887L2V

KRTCAP2 (NM_173852) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	KRTCAP2 (NM_173852) Human Tagged ORF Clone Lentiviral Particle
Symbol:	KRTCAP2
Synonyms:	KCP2
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_173852
ORF Size:	486 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC208887).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 173852.3</u> , <u>NP 776251.1</u>
RefSeq Size:	577 bp
RefSeq ORF:	411 bp
Locus ID:	200185
UniProt ID:	<u>Q8N6L1</u>
Cytogenetics:	1q22
Protein Families:	Transmembrane
MW:	17.4 kDa



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Scrigene KRTCAP2 (NM_173852) Human Tagged ORF Clone Lentiviral Particle – RC208887L2V

Gene Summary:Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a
defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-
pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent
polypeptide chains, the first step in protein N-glycosylation. N-glycosylation occurs
cotranslationally and the complex associates with the Sec61 complex at the channel-forming
translocon complex that mediates protein translocation across the endoplasmic reticulum
(ER). All subunits are required for a maximal enzyme activity (PubMed:22467853). May be
involved in N-glycosylation of APP (amyloid-beta precursor protein). Can modulate gamma-
secretase cleavage of APP by enhancing endoprotelysis of PSEN1 (PubMed:21768116).
[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US