

## Product datasheet for RC208897L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## AGR3 (NM\_176813) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** AGR3 (NM\_176813) Human Tagged ORF Clone Lentiviral Particle

Symbol: AGR3

Synonyms: AG-3; AG3; BCMP11; hAG-3; HAG3; PDIA18

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_176813

ORF Size: 498 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC208897).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 176813.3</u>

 RefSeq Size:
 763 bp

 RefSeq ORF:
 501 bp

 Locus ID:
 155465

 UniProt ID:
 Q8TD06

 Cytogenetics:
 7p21.1

**Protein Families:** Secreted Protein

**MW:** 19.2 kDa







## **Gene Summary:**

This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, a catalytically active thioredoxin domain, and a C-terminal ER-retention sequence. This gene is expressed in ciliated airway epithelial cells and, in mouse, plays a role in ciliary beat frequency in multiciliated cells. This gene is also over-expressed in breast, ovarian, and prostrate cancers. [provided by RefSeq, Dec 2016]