

## **Product datasheet for RC208900**

## GSTM5 (NM 000851) Human Tagged ORF Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** GSTM5 (NM\_000851) Human Tagged ORF Clone

Tag: Myc-DDK
Symbol: GSTM5

**Synonyms:** GSTM5-5; GTM5

Mammalian Cell N

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >RC208900 representing NM\_000851

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC}$ 

GCCGCGATCGCC

AAGTCAGCTACATGGAACAGCAAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RC208900 representing NM\_000851

Red=Cloning site Green=Tags(s)

MPMTLGYWDIRGLAHAIRLLLEYTDSSYVEKKYTLGDAPDYDRSQWLNEKFKLGLDFPNLPYLIDGTHKI TQSNAILRYIARKHNLCGETEEEKIRVDILENQVMDNHMELVRLCYDPDFEKLKPKYLEELPEKLKLYSE FLGKRPWFAGDKITFVDFLAYDVLDMKRIFEPKCLDAFLNLKDFISRFEGLKKISAYMKSSQFLRGLLFG KSATWNSK

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/ja1440-e05.zip">https://cdn.origene.com/chromatograms/ja1440-e05.zip</a>

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 



CTATAGGGCGGCCG	EcoRI GAATTCGT		nHI Kpn I TOOGGTAG	RBS	- GATCTGC		nsus Sgfl	c R	rg -		
ORF	NNN	Mlu I ACG CG T R	T ACG C	Not I GG CCG R IP	Xhol CTC GAG L E			:.Tag : ATC I	TCA S	GAA E	GAG E
GAT CTG GCA GC. D L A A		ATC CTC	GAT TA		g GAT GAC D D	GAC GA		GTT	me I TAA Stop	ACGG	se I COGGOC

<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_000851

ORF Size: 654 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts

of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by

calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

NM 000851.4 RefSeq:

RefSeq Size: 1567 bp RefSeq ORF: 657 bp Locus ID: 2949 UniProt ID: P46439 Cytogenetics: 1p13.3

Domains: GST N, GST C

**Protein Pathways:** Drug metabolism - cytochrome P450, Glutathione metabolism, Metabolism of xenobiotics by

cytochrome P450

MW: 25.5 kDa


**Gene Summary:** Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two

> distinct supergene families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase that belongs to the mu class. The mu class of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs. Diversification of these genes has occurred in regions encoding substrate-binding domains, as well as in tissue expression patterns, to accommodate an increasing number of foreign compounds. [provided by RefSeq,


Jul 2008]



## **Product images:**



Circular map for RC208900



116 — 66 — 45 — 35 — 25 — 18 — 14 — Western blot validation of overexpression lysate (Cat# [LY424493]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208900 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

Coomassie blue staining of purified GSTM5 protein (Cat# [TP308900]). The protein was produced from HEK293T cells transfected with GSTM5 cDNA clone (Cat# RC208900) using MegaTran 2.0 (Cat# [TT210002]).