

Product datasheet for RC208910L4

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

RPL22 (NM_000983) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: RPL22 (NM_000983) Human Tagged Lenti ORF Clone

Tag: mGFP Symbol: RPL22

Synonyms: EAP; HBP15; HBP15/L22; L22

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC208910).

Sequence:

s: Sgfl-Mlul

Restriction Sites: Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_000983

ORF Size: 384 bp



RPL22 (NM_000983) Human Tagged Lenti ORF Clone - RC208910L4

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

1p36.31

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 000983.3</u>

 RefSeq Size:
 2099 bp

 RefSeq ORF:
 387 bp

 Locus ID:
 6146

 UniProt ID:
 P35268

Cytogenetics:

Domains: Ribosomal L22e

Protein Pathways: Ribosome

MW: 15.2 kDa

Gene Summary: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and

a large 60S subunit. Together these subunits are composed of 4 RNA species and


approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 60S subunit. The protein belongs to the L22E family of ribosomal proteins. Its initiating methionine residue is post-translationally removed. The protein can bind specifically to Epstein-Barr virus-encoded RNAs (EBERs) 1 and 2. The mouse protein has been shown to be capable of binding to heparin. Transcript variants utilizing alternative polyA signals exist. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. It was

previously thought that this gene mapped to 3q26 and that it was fused to the acute myeloid leukemia 1 (AML1) gene located at 21q22 in some therapy-related myelodysplastic syndrome patients with 3;21 translocations; however, these fusions actually involve a ribosomal protein L22 pseudogene located at 3q26, and this gene actually maps to 1p36.3-p36.2. [provided by


RefSeq, Jul 2008]



Product images:



Circular map for RC208910L4



Double digestion of RC208910L4 using Sgfl and Mlul $\,$