

Product datasheet for **RC208917**

MMP14 (NM_004995) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	MMP14 (NM_004995) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	MMP14
Synonyms:	MMP-14; MMP-X1; MT-MMP; MT-MMP 1; MT1-MMP; MT1MMP; MTMMP1; WNCHRS
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Cloning Scheme:


ACCN: NM_004995

ORF Size: 1746 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [NM_004995.4](#)


RefSeq Size: 3558 bp

RefSeq ORF: 1749 bp


Locus ID: 4323

UniProt ID:	P50281
Cytogenetics:	14q11.2
Domains:	hemopexin, Peptidase_M10, ZnMc
Protein Families:	Druggable Genome, Protease, Transmembrane
Protein Pathways:	GnRH signaling pathway
MW:	65.89 kDa
Gene Summary:	<p>Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the protein encoded by this gene is a member of the membrane-type MMP (MT-MMP) subfamily; each member of this subfamily contains a potential transmembrane domain suggesting that these proteins are expressed at the cell surface rather than secreted. This protein activates MMP2 protein, and this activity may be involved in tumor invasion. [provided by RefSeq, Jul 2008]</p>


Product images:



Circular map for RC208917



Western blot validation of overexpression lysate (Cat# [LY401554]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC208917 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified MMP14 protein (Cat# [TP308917]). The protein was produced from HEK293T cells transfected with MMP14 cDNA clone (Cat# RC208917) using MegaTran 2.0 (Cat# [TT210002]).