

Product datasheet for RC208919L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ICAM3 (NM_002162) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ICAM3 (NM_002162) Human Tagged ORF Clone Lentiviral Particle

Symbol: ICAM3

Synonyms: CD50; CDW50; ICAM-R

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_002162

 ORF Size:
 1641 bp

ORF Nucleotide Sequence:

The ORF insert of this clone is exactly the same as(RC208919).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002162.2

 RefSeq Size:
 1796 bp

 RefSeq ORF:
 1644 bp

 Locus ID:
 3385

 UniProt ID:
 P32942

 Cytogenetics:
 19p13.2

Domains: ig, IG, ICAM_N

Protein Families: ES Cell Differentiation/IPS, Transmembrane





ICAM3 (NM_002162) Human Tagged ORF Clone Lentiviral Particle - RC208919L3V

Protein Pathways: Cell adhesion molecules (CAMs)

MW: 59.5 kDa

Gene Summary: The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM)

family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9

immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This protein is constitutively and abundantly expressed by all leucocytes and may be the most important ligand for LFA-1 in the initiation of the immune response. It functions not only as an adhesion molecule, but also as a potent signalling molecule. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2016]