

## Product datasheet for RC208922L2V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# CDA (NM 001785) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type: Lentiviral Particles** 

**Product Name:** CDA (NM\_001785) Human Tagged ORF Clone Lentiviral Particle

Symbol: CDA CDD Synonyms: **Mammalian Cell** 

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

mGFP Tag:

NM 001785 ACCN:

**ORF Size:** 438 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC208922).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001785.1, NP 001776.1

RefSeq Size: 892 bp RefSeq ORF: 441 bp Locus ID: 978 **UniProt ID:** P32320

Cytogenetics: 1p36.12

**Domains:** dCMP\_cyt\_deam

**Protein Families:** Stem cell - Pluripotency





### CDA (NM\_001785) Human Tagged ORF Clone Lentiviral Particle - RC208922L2V

**Protein Pathways:** Drug metabolism - other enzymes, Metabolic pathways, Pyrimidine metabolism

**MW:** 16 kDa

**Gene Summary:** This gene encodes an enzyme involved in pyrimidine salvaging. The encoded protein forms a

homotetramer that catalyzes the irreversible hydrolytic deamination of cytidine and deoxycytidine to uridine and deoxyuridine, respectively. It is one of several deaminases responsible for maintaining the cellular pyrimidine pool. Mutations in this gene are associated with decreased sensitivity to the cytosine nucleoside analogue cytosine

arabinoside used in the treatment of certain childhood leukemias. [provided by RefSeq, Jul  $\,$ 

2008]