

Product datasheet for RC208935L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TC10 (RHOQ) (NM_012249) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TC10 (RHOQ) (NM_012249) Human Tagged ORF Clone Lentiviral Particle

Symbol: TC10

Synonyms: ARHQ; HEL-S-42; RASL7A; TC10; TC10A

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_012249

ORF Size: 615 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC208935).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 012249.2

 RefSeq Size:
 4542 bp

 RefSeq ORF:
 618 bp

 Locus ID:
 23433

 UniProt ID:
 P17081

 Cytogenetics:
 2p21

, ,

Domains: ras, RAS, RHO, RAB

Protein Pathways: Insulin signaling pathway





ORÏGENE

MW: 22.7 kDa

Gene Summary:

This gene encodes a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. The encoded protein is an important signalling protein for sarcomere assembly and has been shown to play a significant role in the exocytosis of the solute carrier family 2, facilitated glucose transporter member 4 and other proteins, possibly acting as the signal that turns on the membrane fusion machinery. Three related pseudogene have been identified on chromosomes 2 and 14. [provided by RefSeq, Aug 2011]