

Product datasheet for RC209001L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PSMB9 (NM 002800) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: PSMB9 (NM_002800) Human Tagged ORF Clone Lentiviral Particle

Symbol: PSMB9

Synonyms: beta1i; LMP2; PRAAS3; PSMB6i; RING12

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002800

ORF Size: 657 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209001).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002800.4

 RefSeq Size:
 1048 bp

 RefSeq ORF:
 660 bp

 Locus ID:
 5698

 UniProt ID:
 P28065

 Cytogenetics:
 6p21.32

Domains: proteasome

Protein Families: Druggable Genome, Protease





Protein Pathways: Proteasome

MW: 23.3 kDa

Gene Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S

core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is induced by gamma interferon and this

gene product replaces catalytic subunit 1 (proteasome beta 6 subunit) in the

immunoproteasome. Proteolytic processing is required to generate a mature subunit.

[provided by RefSeq, Mar 2010]