

Product datasheet for RC209021L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ADCY6 (NM_015270) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ADCY6 (NM_015270) Human Tagged ORF Clone Lentiviral Particle

Symbol: ADCY6

Synonyms: AC6; LCCS8

Mammalian Cell

11

Puromycin

Selection: Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_015270 **ORF Size:** 3504 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209021).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 015270.3

RefSeq Size:6583 bpRefSeq ORF:3507 bp

Locus ID: 112

 UniProt ID:
 O43306

 Cytogenetics:
 12q13.12

Domains: CYCc

Protein Families: Druggable Genome, Transmembrane





ADCY6 (NM_015270) Human Tagged ORF Clone Lentiviral Particle - RC209021L4V

Protein Pathways: Chemokine signaling pathway, Dilated cardiomyopathy, Gap junction, GnRH signaling

pathway, Melanogenesis, Oocyte meiosis, Progesterone-mediated oocyte maturation, Purine

metabolism, Taste transduction, Vascular smooth muscle contraction

MW: 130.6 kDa

Gene Summary: This gene encodes a member of the adenylyl cyclase family of proteins, which are required

for the synthesis of cyclic AMP. All members of this family have an intracellular N-terminus, a tandem repeat of six transmembrane domains separated by a cytoplasmic loop, and a C-terminal cytoplasmic domain. The two cytoplasmic regions bind ATP and form the catalytic core of the protein. Adenylyl cyclases are important effectors of transmembrane signaling pathways and are regulated by the activity of G protein coupled receptor signaling. This protein belongs to a small subclass of adenylyl cyclase proteins that are functionally related and are inhibited by protein kinase A, calcium ions and nitric oxide. A mutation in this gene is

associated with arthrogryposis multiplex congenita. [provided by RefSeq, May 2015]