

Product datasheet for **RC209069L1V**

Pinin (PNN) (NM_002687) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Pinin (PNN) (NM_002687) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Pinin
Synonyms:	DRS; DRSP; memA; SDK3
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_002687
ORF Size:	2151 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC209069).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_002687.3
RefSeq Size:	3606 bp
RefSeq ORF:	2154 bp
Locus ID:	5411
UniProt ID:	Q9H307
Cytogenetics:	14q21.1
Domains:	pinin_SDK_memA, pinin_SDK_N
Protein Families:	Stem cell - Pluripotency, Transcription Factors



[View online »](#)

MW: 81.6 kDa

Gene Summary: Transcriptional activator binding to the E-box 1 core sequence of the E-cadherin promoter gene; the core-binding sequence is 5'CAGGTG-3'. Capable of reversing CTBP1-mediated transcription repression. Auxiliary component of the splicing-dependent multiprotein exon junction complex (EJC) deposited at splice junction on mRNAs. The EJC is a dynamic structure consisting of core proteins and several peripheral nuclear and cytoplasmic associated factors that join the complex only transiently either during EJC assembly or during subsequent mRNA metabolism. Participates in the regulation of alternative pre-mRNA splicing. Associates to spliced mRNA within 60 nt upstream of the 5'-splice sites. Component of the PSAP complex which binds RNA in a sequence-independent manner and is proposed to be recruited to the EJC prior to or during the splicing process and to regulate specific excision of introns in specific transcription subsets. Involved in the establishment and maintenance of epithelia cell-cell adhesion. Potential tumor suppressor for renal cell carcinoma.[UniProtKB/Swiss-Prot Function]