

Product datasheet for RC209110L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ALKBH5 (NM_017758) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ALKBH5 (NM_017758) Human Tagged ORF Clone Lentiviral Particle

Symbol: ALKBH5

Synonyms: ABH5; OFOXD; OFOXD1

Mammalian Cell None

Selection:

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_017758 **ORF Size:** 1374 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC209110).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 017758.2, NP 060228.2

 RefSeq Size:
 3449 bp

 RefSeq ORF:
 1185 bp

 Locus ID:
 54890

 UniProt ID:
 Q6P6C2

 Cytogenetics:
 17p11.2

Domains: 20G-Fell_Oxy

MW: 51.4 kDa







Gene Summary:

Dioxygenase that demethylates RNA by oxidative demethylation: specifically demethylates N(6)-methyladenosine (m6A) RNA, the most prevalent internal modification of messenger RNA (mRNA) in higher eukaryotes (PubMed:23177736, PubMed:24489119, PubMed:24616105, PubMed:24778178). Can also demethylate N(6)-methyladenosine in single-stranded DNA (in vitro) (PubMed:24616105). Requires molecular oxygen, alpha-ketoglutarate and iron (PubMed:21264265, PubMed:23177736, PubMed:24489119, PubMed:24616105, PubMed:24778178). Demethylation of m6A mRNA affects mRNA processing and export (PubMed:23177736). Required for the late meiotic and haploid phases of spermatogenesis by mediating m6A demethylation in spermatocytes and round spermatids: m6A demethylation of target transcripts is required for correct splicing and the production of longer 3' UTR mRNAs in male germ cells (By similarity).[UniProtKB/Swiss-Prot Function]