

## Product datasheet for RC209120L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PLC delta 3 (PLCD3) (NM 133373) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** PLC delta 3 (PLCD3) (NM 133373) Human Tagged ORF Clone Lentiviral Particle

Symbol:

PLC-delta-3 Synonyms:

**Mammalian Cell** 

Selection:

ACCN:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 133373

**ORF Size:** 2367 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC209120).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 133373.3, NP 588614.1

RefSeq Size: 3477 bp RefSeq ORF: 2370 bp Locus ID: 113026 **UniProt ID:** Q8N3E9 Cytogenetics: 17q21.31

**Domains:** C2, PI-PLC-X, PI-PLC-Y, PH

**Protein Families:** Druggable Genome





## PLC delta 3 (PLCD3) (NM\_133373) Human Tagged ORF Clone Lentiviral Particle - RC209120L1V

**Protein Pathways:** Calcium signaling pathway, Inositol phosphate metabolism, Metabolic pathways,

Phosphatidylinositol signaling system

**MW:** 89.1 kDa

**Gene Summary:** This gene encodes a member of the phospholipase C family, which catalyze the hydrolysis of

phosphatidylinositol 4,5-bisphosphate to generate the second messengers diacylglycerol and

inositol 1,4,5-trisphosphate (IP3). Diacylglycerol and IP3 mediate a variety of cellular

responses to extracellular stimuli by inducing protein kinase C and increasing cytosolic Ca(2+) concentrations. This enzyme localizes to the plasma membrane and requires calcium for activation. Its activity is inhibited by spermine, sphingosine, and several phospholipids.

[provided by RefSeq, Jul 2008]